arXiv:1212.0351v1 [cond-mat.str-el] 3 Dec 2012

Condensate-induced transitions and critical spin chains

Teresia Månsson,[1] Ville Lahtinen,[2,3] Juha Suorsa,[2] and Eddy Ardonne[2,4]
[1]*Department of Theoretical Physics, School of Engineering Sciences,*
Royal Institute of Technology (KTH), Roslagstullsbacken 21, SE-106 91 Stockholm, Sweden
[2]*Nordita, Royal Institute of Technology and Stockholm University,*
Roslagstullsbacken 23, SE-106 91 Stockholm, Sweden
[3]*University of Amsterdam, Institute of Physics, Science Park 904, 1090 GL Amsterdam, The Netherlands*
[4]*Department of Physics, Stockholm University, AlbaNova University Center, SE-106 91 Stockholm, Sweden*
(Dated: December 4, 2012)

We show that the framework of condensate-induced transitions between two-dimensional topological phases can be used to relate one-dimensional spin models at their critical points. In particular, we show that two well-known spin chains, namely the XY chain and the transverse field Ising chain with only next-nearest-neighbor interactions, differ at their critical points only by a boundary term and can be related via an exact mapping. The boundary term constrains the symmetry sectors of the transverse field Ising chain, reducing the number of primary fields in the conformal field theory that describes its critical behavior. We argue that the reduction of the field content is equivalent to the confinement of a set of primary fields, in analogy to the confinement of quasiparticles resulting from a condensation of a boson in a topological phase. To provide evidence for the generality of the framework we apply it to the XY chain with only next-nearest-neighbor interactions. In the presence of a confining boundary term, this XY chain can be mapped to a spin chain with the $u(1)_2 \times u(1)_2$ critical behavior predicted by the condensation framework.

An essential characteristic of topologically ordered phases is the absence of local order parameters [1]. While this property ensures a degree of stability, which is of potential benefit in quantum information technologies, it also makes it necessary to develop new approaches for the study of phase transitions in these systems. As the low-energy degrees of freedom are anyonic quasiparticles specific to a particular topological phase, it should be possible to understand transitions between the phases in terms of their collective behavior. One successful framework is that of condensate-induced transitions [2], where the condensation of a bosonic particle induces confinement of a set of quasiparticles, thereby causing a transition to a new phase. Topological phase transitions in various systems, including the quantum Hall hierarchies [3–7], interacting anyons [8–10] and Levin-Wen models [11, 12], can be related to this mechanism. It is well known that 2+1-dimensional topological phases are intimately related to two-dimensional conformal field theories (CFT) [13, 14]. In particular, the topological charges carried by the bulk quasiparticles are in one-to-one correspondence with the primary fields of a CFT, which also describes the gapless edge modes of the topological phase. It is then natural to ask whether the condensate-induced transitions have a counterpart in critical one-dimensional systems that are described by CFT's.

In this Letter, we answer this question positively and show that the framework of condensate-induced transitions applies in the context of critical one-dimensional spin chains. We demonstrate this by considering two canonical spin chains: the XY chain, introduced and solved by Lieb, Schultz and Mattis [15], and the transverse field Ising chain [16, 17], in which only the next-nearest-neighbor spins interact. The criticality of these chains is described by the $u(1)_4$ and the Ising × Ising CFT's, respectively, which are related by the condensation framework [2] when viewed as low-energy theories of topological phases. We show that for periodic boundary conditions, these two spin chains differ at their critical points only by a boundary term that constrains the symmetry sectors of the transverse field Ising chain. We argue that these constraints, in effect, confine those CFT primary fields which correspond to bulk quasiparticles confined due to condensation. To demonstrate the generality of such relations between critical spin chains, we apply a similar confining boundary term to an XY chain with only next-nearest neighbor interactions and derive a critical microscopic spin chain with the predicted $u(1)_2 \times u(1)_2$ CFT behavior. The framework of condensate-induced transitions thus offers a powerful general framework for relating known critical spin chains and deriving novel ones from them.

The transverse field Ising model – The critical transverse field Ising (TFI) chains that we consider are given by the Hamiltonian

$$H^n_{\mathrm{TFI}} = \sum_{i=0}^{L-1} \sigma^z_i + \sigma^x_i \sigma^x_{i+n} \, , \qquad (1)$$

where we assume periodic boundary conditions (PBC), $\sigma^\alpha_L = \sigma^\alpha_0$. For $n = 1$ we obtain the critical nearest-neighbor chain. This model can be solved via a Jordan-Wigner transformation, which transforms the Pauli operators σ^α_i into fermionic operators $c^\dagger_i$ and c_i [16, 17]. The operators $c^\dagger_i$ involve strings of Pauli operators, which we choose to start at site zero and end at site i. The parity of the number of fermions F is a symmetry of the Hamiltonian, and the corresponding symmetry operator is given by $\mathcal{P} = \prod_{i=0}^{L-1} \sigma_i^z = (-1)^F$. In terms of the fermions, the Hamiltonian reads

$$H^1_{\mathrm{TFI}} = \sum_{i=0}^{L-1}(2c_i^\dagger c_i - 1) + \sum_{i=0}^{L-2}(c_i - c_i^\dagger)(c_{i+1} + c_{i+1}^\dagger) - (-1)^F (c_{L-1} - c_{L-1}^\dagger)(c_0 + c_0^\dagger)\,. \tag{2}$$

The boundary conditions thus depend on the parity $\mathcal{P}$: for odd F, one has $c_L \equiv c_0$, while for even F, one has $c_L \equiv -c_0$. In momentum space the Hamiltonian H^1_{TFI} takes the form (after a Bogoliubov transformation)

$$H^1_{\mathrm{TFI}} = -4 \sum_k \left|\sin\left(\frac{\pi k}{L}\right)\right| \left(c_k^\dagger c_k - \frac{1}{2}\right) , \tag{3}$$

where the $c_k^\dagger$ create fermions with momentum k. Due to the parity-dependent boundary conditions, these momenta take integer values for odd F, and half-integer values for even F. The CFT describing the criticality of the TFI chain is the Ising CFT with central charge $c = 1/2$. This CFT has three primary fields, 1, σ and ψ, with scaling dimensions $h_1 = 0$, $h_\sigma = 1/16$ and $h_\psi = 1/2$. The even fermion sector with $\mathcal{P} = 1$ contains the states corresponding to the primary fields $\mathbf{1}$ and ψ, while the odd fermion sector $\mathcal{P} = -1$ contains the states corresponding to the primary field σ. We refer to [18] for more details on CFT.

Setting $n = 2$ in (1), the model has only next-nearest-neighbor interactions. The chain decouples into two TFI chains on the even (e) and the odd (o) sites, each of which can be solved in the same way as the ordinary transverse field Ising model above. The fermion parity is conserved independently for the even and odd sites with the corresponding symmetry operators now given by $\mathcal{P}_e = \prod_j \sigma_{2j}^z$ and $\mathcal{P}_o = \prod_j \sigma_{2j+1}^z$. The CFT describing the critical behavior is the direct product of two Ising CFT's (denoted as Ising^2), with total central charge $c = 1$. The correspondence between the symmetry sectors of H^2_{TFI} and the primary fields of the Ising^2 CFT is shown in Table I.

Sectors of H^2_{TFI}

($\mathcal{P}_e$,$\mathcal{P}_o$)	Ising^2 fields
$(1,1)$	$(1,1)$,$(1,\psi)$,$(\psi,1)$,(ψ,ψ)
$(1,-1)$	$(1,\sigma)$, (ψ,σ)
$(-1,1)$	$(\sigma,1)$, (σ,ψ)
$(-1,-1)$	(σ,σ)

Sectors of H^1_{XY}

$\mathcal{T}^z$	$u(1)_4$ fields
1	$\tilde{1}$, $\tilde{\psi}$
-1	λ, $\bar{\lambda}$

TABLE I. *Left:* The correspondence between the symmetry sectors of H^2_{TFI} and the sectors of the Ising^2 CFT. *Right:* The correspondence between the symmetry sectors of H^1_{XY} and the sectors of the $u(1)_4$ CFT.

The XY spin chain – The second spin chain we consider is the critical XY chain. Its Hamiltonian is given by

$$H^n_{\mathrm{XY}} = \sum_{i=0}^{L-1} \tau_i^x \tau_{i+n}^x + \tau_i^y \tau_{i+n}^y \ , \tag{4}$$

where τ_i^α denote the Pauli matrices acting on the site i. We assume periodic boundary conditions and even L. As above, the case $n = 1$ ($n = 2$) corresponds to a model with only nearest-neighbor (next-nearest-neighbor) interactions. The XY spin chain too can be brought into a diagonal form via a Jordan-Wigner transformation [15]. The operator $\mathcal{T}^z = \prod_i \tau^z$ commutes with H^n_{XY}. It again corresponds to the fermion parity and determines the boundary conditions. In momentum space, the nearest-neighbor XY chain is diagonal:

$$H^1_{\mathrm{XY}} = 4 \sum_k \cos\left(\frac{2\pi k}{L}\right) \left(c_k^\dagger c_k - \frac{1}{2}\right) \ . \tag{5}$$

The spectrum of the XY chain at the critical point can be described in terms of the $u(1)_4$ CFT of a chiral boson [18]. This CFT has central charge $c = 1$ and contains four primary fields $\tilde{1}$, λ, $\bar{\lambda}$ and $\tilde{\psi}$ with scaling dimensions $h_{\tilde{1}} = 0$, $h_\lambda = h_{\bar{\lambda}} = 1/8$ and $h_{\tilde{\psi}} = 1/2$.

As illustrated in Table I, states descending from the vacuum $\tilde{1}$ and fermion $\tilde{\psi}$ primary fields appear in the even parity sector $\mathcal{T}^z = 1$ while the states descending from the λ, $\bar{\lambda}$ fields appear in the odd parity sector $\mathcal{T}^z = -1$. For a chain of *even* length the operators $\mathcal{T}^x = \prod_{i=0}^{L-1} \tau_i^x$ and $\mathcal{T}^y = \prod_{i=0}^{L-1} \tau_i^y$ commute with the Hamiltonian Eq. (4), with each other and with $\mathcal{T}^z$. So for even L, the sectors $\mathcal{T}^z = \pm 1$ each split into two sectors, which are degenerate in the case $\mathcal{T}^z = -1$.

We will show below that when a suitable boundary term H^B_{TFI} is introduced, $H^2_{\mathrm{TFI}} + H^B_{\mathrm{TFI}}$ can be mapped exactly to H^1_{XY}. The boundary terms we consider are non-local in the sense that they change the boundary conditions, depending on the symmetry sector, similarly to the boundary term in the $n = 1$ TFI chain in Eq. (2). To assign a physical meaning to the boundary term, we set the spin chains briefly aside and review the notion of condensate-induced transitions between gapped topological phases [2].

The condensation framework – In topological field theories, condensate-induced transitions proceed in three steps that we outline in the context of the Ising^2 theory. The labels 1, ψ and σ of an Ising theory should be understood as topological quantum numbers of the quasi-particle excitations with the following 'tensor product', or fusion rules:

$$\psi \times \sigma = \sigma, \quad \psi \times \psi = 1, \quad \sigma \times \sigma = 1 + \psi\,. \tag{6}$$

Fusion is symmetric ($a \times b = b \times a$), distributive ($(a+b) \times c = a \times c + b \times c$), associative ($a \times (b \times c) = (a \times b) \times c$) and all particles fuse trivially with the vacuum label ($a \times 1 = a$).

The fusion rules of the Ising^2 theory follow directly from the fusion rules of each Ising theory.

In our example, the condensation is effected by identifying the boson (ψ,ψ) with the vacuum label $(1,1)$. By definition, the vacuum should have trivial fusion and statistics with all the other particles. Thus, in order that (ψ,ψ) behaves like the vacuum, the first step is to identify all particles a and b that are related by fusion with it. That is, if $a\times(\psi,\psi)=b$, then we set $a=b$. We arrive at a new set of particle types $\tilde{1}=(1,1)=(\psi,\psi)$, $\tilde{\psi}=(1,\psi)=(\psi,1)$, $\tilde{\sigma}_1=(\sigma,1)=(\sigma,\psi)$ and $\tilde{\sigma}_2=(1,\sigma)=(\psi,\sigma)$, while the particle (σ,σ) remains unaffected at this step. In the second step, one demands that all particles should have trivial statistics with the new vacuum. This is equivalent to demanding that identified particles with unequal conformal weights are confined. Since $h_{(\sigma,1)}=1/16$, but $h_{(\sigma,\psi)}=9/16$, the particles $\tilde{\sigma}_1$ and $\tilde{\sigma}_2$ are eliminated from the particle content of the condensed phase. At the third and final step, one verifies that the remaining particles have consistent fusion rules, i.e. the new vacuum is unique and the fusion algebra closes. We find this not to be the case, because $(\sigma,\sigma)\times(\sigma,\sigma)=(1,1)+(1,\psi)+(\psi,1)+(\psi,\psi)=2\cdot\tilde{1}+2\cdot\tilde{\psi}$, which means that (σ,σ) must branch into new particle types. The uniqueness of the vacuum can be satisfied if one replaces (σ,σ) by $\lambda+\bar{\lambda}$ and demands that these particles satisfy the new fusion rules

$$\begin{gathered}\tilde{\psi}\times\lambda=\bar{\lambda},\quad \tilde{\psi}\times\bar{\lambda}=\lambda,\quad \tilde{\psi}\times\tilde{\psi}=\tilde{1}\,,\\ \lambda\times\bar{\lambda}=\tilde{1},\quad \lambda\times\lambda=\bar{\lambda}\times\bar{\lambda}=\tilde{\psi}\,.\end{gathered}\tag{7}$$

These are the fusion rules of the $u(1)_4$ theory, which is the theory obtained from the Ising^2 theory via a condensation transition [2].

Condensation and critical spin chains – As the topological quantum numbers of the quasiparticles are in correspondence with the primary fields of the CFT that describes the edge of the topological phase, the condensation mechanism suggests a similar connection between the Ising^2 and $u(1)_4$ CFT's that describe the critical behavior of the H^2_{TFI} and H^1_{XY} chains. To induce the same effect as condensing the (ψ,ψ) field we eliminate the sectors that contain the confined $\tilde{\sigma}_1$ and $\tilde{\sigma}_2$ particles. This can be achieved by the boundary term

$$H^B_{\mathrm{TFI}}=\left(\mathcal{P}_o-\mathbf{1}\right)\sigma^x_{L-2}\sigma^x_0+\left(\mathcal{P}_e-\mathbf{1}\right)\sigma^x_{L-1}\sigma^x_1\,,\tag{8}$$

that couples the chains on the even and odd sites. Namely, the odd parity sector of one chain forces anti-periodic boundary conditions in the other, effectively changing the parity sector of the latter. It follows that the overall parity is even. Because the parity determines the sectors of the CFT description, we find that the sectors $\tilde{\sigma}_1$ and $\tilde{\sigma}_2$, with odd overall parity, are eliminated, as shown in table II.

Condensed phase $H^2_{\mathrm{TFI}}+H^B_{\mathrm{TFI}}=H^1_{\mathrm{XY}}$

($\mathcal{P}_e\mathcal{P}_o$,$\mathcal{P}_o\mathcal{P}_e$)	Ising^2 fields	$\mathcal{T}^z$	$u(1)_4$ fields
$(1,1)$	$(1,1)$,$(1,\psi)$,$(\psi,1)$,(ψ,ψ)	1	$1,\tilde{\psi}$
$(-1,-1)$	(σ,σ)	-1	$\lambda,\bar{\lambda}$
$(-1,-1)$	(σ,σ)	-1	$\lambda,\bar{\lambda}$
$(1,1)$	$(1,1)$,$(1,\psi)$,$(\psi,1)$,(ψ,ψ)	1	$1,\tilde{\psi}$

TABLE II. When the boundary term H^B_{TFI} is added to H^2_{TFI}, the boundary conditions on each chain depend on the parity sectors of both chains. The sectors of $H^2_{\mathrm{TFI}}+H^B_{\mathrm{TFI}}$ are thus labeled by ($\mathcal{P}_e\mathcal{P}_o$,$\mathcal{P}_o\mathcal{P}_e$), which reduces the number of sectors from four to two. Because ($\mathcal{P}_e$,$\mathcal{P}_o$)=($\mathcal{T}^x$,$\mathcal{T}^y$), the two surviving sectors map to the parity sectors of H^1_{XY} labeled by $\mathcal{T}^z=\mathcal{T}^x\mathcal{T}^y$. As shown in Fig. 1, the states in these sectors can be uniquely associated with the identified/branched states predicted by the condensation mechanism.

An explicit relation between the next-nearest-neighbor TFI and the XY chain can be obtained by a canonical transformation inspired by the one used in [19] for open chains. The non-local and site-dependent transformation

$$\begin{aligned}\sigma^z_{2j}&=\tau^y_{2j}\tau^y_{2j+1} & \sigma^z_{2j+1}&=\tau^x_{2j}\tau^x_{2j+1}\\ \sigma^x_{2j}&=\prod_{i\leq 2j}\tau^x_i & \sigma^x_{2j+1}&=\prod_{i\geq 2j+1}\tau^y_i\,,\end{aligned}\tag{9}$$

on the spin variables in the TFI gives

$$H^2_{\mathrm{TFI}}+H^B_{\mathrm{TFI}}=H^1_{\mathrm{XY}}\,.\tag{10}$$

This establishes a microscopic relation between the two chains and is a central result of this work.

To justify that H^B_{TFI} indeed induces an analogue of a condensation transition, we show that the spectra of the surviving sectors encode also the two other defining features of the mechanism: the identification and the splitting of particles (in addition to the confinement of the $\tilde{\sigma}_1$ and $\tilde{\sigma}_2$ sectors). To show this, we realize that (9) gives $\mathcal{P}_e=\mathcal{T}^y$ and $\mathcal{P}_o=\mathcal{T}^x$, i.e. the sectors of H^2_{TFI} chain correspond exactly to sectors of the H^1_{XY} chain. For $\mathcal{P}_e=\mathcal{P}_o=1$, the boundary term Eq. (8) is zero, so in this sector, H^2_{TFI} and H^1_{XY} are equal. In the other sectors, the energies differ, while the momenta of the states will in general be shifted, due to the change in boundary conditions. Fig. 1 shows that when the spectra of these models are plotted on top of each other and the individual states are labeled according to the CFT predictions, we confirm the condensation picture. Namely, the identification brought about by the condensation is confirmed by noting that (ψ,ψ) states map to the new vacuum $\tilde{1}$ states, while states descending from $(1,\psi)$ and $(\psi,1)$ map to the new fermion $\tilde{\psi}$ states. The branching $(\sigma,\sigma)\to\lambda+\bar{\lambda}$ manifests itself as the two-fold degeneracy of all states corresponding to these fields.

Deriving critical spin models – The condensation mechanism underlying the boundary term and the subsequent mapping between the $n=2$ TFI and $n=1$ XY chains suggests that similar mappings should also hold



FIG. 1. The low-part of the rescaled energy spectra of the $n = 2$ TFI chain (crosses) and the XY chain (squares) for $L = 20$. The spectra are symmetric under $K \to -K$. The brown crosses correspond to the sectors of the $n = 2$ TFI chain which become confined after the condensation. The blue, red and black crosses correspond to the $(1,1)$, $(\psi,1)$ and (σ,σ) sectors, respectively. The blue, red and black squares correspond to the $\tilde{1}$, $\tilde{\psi}$ and $\lambda/\bar{\lambda}$ sectors of the XY chain. The dashed lines correspond to the energies predicted by CFT, namely $E = h + \bar{h} + k$, where $h, \bar{h}$ are the left and right conformal dimensions of the primary fields and k is an integer. In the XY chain, $h + \bar{h}$ takes the values $0, 1/8, 1$.

between other critical spin chains whose CFT descriptions can be related by the condensation picture. We employ this insight to derive a critical spin chain described by the $u(1)_2 \times u(1)_2$ CFT.

We start with the critical $n = 2$ XY chain, which is described by the $u(1)_4 \times u(1)_4$ CFT, containing sixteen primary fields (compare to the $n = 2$ TFI chain). Without going through the details of the condensation picture, we note that this CFT contains a bosonic field, $(\tilde{\psi}, \tilde{\psi})$. Condensation of this boson leads to a pairwise identification of the 16 fields. Four of the eight resulting pairs turn out to be confined, while none of the fields have to be split. In the end, one is left with a theory with four fields, whose scaling dimensions and fusion rules match those of the $u(1)_2 \times u(1)_2$ CFT.

To induce the counterpart of this transition in $H^2_{\rm XY}$, we again remove the odd fermion parity sectors that contain the confined fields, by introducing the boundary term

$$\begin{aligned} H^B_{\rm XY} = \big(\mathcal{T}^z_o - \mathbf{1}\big)\big(\tau^x_{L-2}\tau^x_0 + \tau^y_{L-2}\tau^y_0\big) + \\ \big(\mathcal{T}^z_e - \mathbf{1}\big)\big(\tau^x_{L-1}\tau^x_1 + \tau^y_{L-1}\tau^y_1\big). \end{aligned} \tag{11}$$

Using the non-local transformation

$$\begin{aligned} \tau^z_{2j} &= \rho^y_{2j}\rho^y_{2j+1} & \tau^z_{2j+1} &= \rho^x_{2j}\rho^x_{2j+1} \\ \tau^x_{2j} &= \Big(\prod_{i<2j}\rho^x_i\Big)\rho^z_{2j}\rho^y_{2j+1} & \tau^x_{2j+1} &= \prod_{i\geq 2j+1}\rho^y_i \\ \tau^y_{2j} &= \prod_{i\leq 2j}\rho^x_i & \tau^y_{2j+1} &= \rho^z_{2j}\rho^z_{2j+1}\Big(\prod_{i>2j+1}\rho^y_i\Big)\ , \end{aligned} \tag{12}$$

where the label j takes the values $j = 0, 1, \ldots, L/2 - 1$ and ρ^α are new Pauli matrices, we arrive at the local and translationally invariant Hamiltonian $H^2_{\rm XY} + H^B_{\rm XY} = \hat{H}$, where $\hat{H}$ is given by

$$\begin{aligned} \hat{H} = \sum_{j=0}^{L/2-1} \Big(&\rho^x_{2j}\rho^z_{2j+1}\rho^z_{2j+2}\rho^x_{2j+3} + \rho^x_{2j+1}\rho^x_{2j+2} \\ &+ \rho^y_{2j}\rho^z_{2j+1}\rho^z_{2j+2}\rho^y_{2j+3} + \rho^y_{2j+1}\rho^y_{2j+2}\Big)\ . \end{aligned} \tag{13}$$

The resulting chain $\hat{H}$ can be solved by means of a Jordan-Wigner transformation, and we have verified that the critical behavior is indeed described by the $u(1)_2 \times u(1)_2$ CFT, as predicted by the condensation framework. Although both the starting and resulting CFT's both have a direct product structure, the condensation mixes the two $u(1)_4$ factors describing the critical behavior of the $n = 2$ XY chain. We provide more details on the condensation transition of the $n = 2$ XY chain in a forthcoming publication [20].

Conclusions – We have shown that two copies of the critical transverse field Ising chain coupled by a boundary term can be related to the critical XY chain via an exact mapping. The key observation is that the coupling through the boundary term constrains the allowed symmetry sectors of the two TFI chains in a manner that is analogous to the confinement induced by condensation of a quasiparticle in a 2D topological phase. Additionally, we have argued for the generality of the framework by applying a similar constraint to two copies of the critical XY chain to obtain a novel critical spin model with a $u(1)_2 \times u(1)_2$ CFT description, as predicted by the corresponding condensate-induced transition.

We expect that the applicability of this framework extends beyond our examples. Including in a critical spin chain a boundary term which eliminates the symmetry sectors that contain the confined primary fields, the critical behavior of the resulting system can be predicted from the condensation picture. In the examples considered, we found non-local transformations of the spin operators leading to manifestly local and translation-invariant Hamiltonians. A more general analysis, along the lines of Ref. [21], using methods of boundary conformal field theory should be possible.

Acknowledgements – EA thanks T. Quella for stimulating discussions. The research of TM is supported by the Swedish science research council.

---

[1] X.G. Wen, *Topological orders and Edge excitations in FQH states*, Adv. Phys. **44**, 405 (1995).

[2] F.A. Bais, J.K. Slingerland, *Condensate-induced transitions between topologically ordered phases*, Phys. Rev. B **79**, 045316 (2009).

[3] B.I. Halperin, *Theory of the quantized Hall conductance*, Helv. Phys. Acta **56** 75 (1983).

[4] F. Haldane, *Fractional quantization of the Hall effect: a hierarchy of incompressible quantum fluid states*, Phys. Rev. Lett. **51**, 605 (1983).

[5] J. Suorsa, S. Viefers and T.H. Hansson, *A general approach to quantum Hall hierarchies*, New J. Phys. **13**, 075006 (2011).

[6] P. Bonderson, *Hierarchical Nature of the Quantum Hall Effects*, Phys. Rev. Lett. **108**, 066806 (2012).

[7] M. Barkeshli and X.-G. Wen, *Anyon Condensation and Continuous Topological Phase Transitions in Non-Abelian Fractional Quantum Hall States*, Phys. Rev. Lett. **105**, 216804 (2010).

[8] C. Gils, E. Ardonne, S. Trebst, A. W. W. Ludwig, M. Troyer and Z. Wang, *Collective States of Interacting Anyons, Edge States, and the Nucleation of Topological Liquids*, Phys. Rev. Lett. **103**, 070401 (2009).

[9] A. W.W. Ludwig, D. Poilblanc, S. Trebst and M. Troyer, *Two-dimensional quantum liquids from interacting non-Abelian anyons*, New J. Phys. **13**, 045014 (2011).

[10] V. Lahtinen, A. W. W. Ludwig, J. K. Pachos and Simon Trebst, *Topological liquid nucleation induced by vortex-vortex interactions in Kitaev's honeycomb model*, Phys. Rev. B **86**, 075115 (2012).

[11] M.A. Levin and X.-G. Wen, *String-net condensation: A physical mechanism for topological phases*, Phys. Rev. B 71, 045110 (2005).

[12] F. J. Burnell, S. H. Simon and J. K. Slingerland, *Condensation of achiral simple currents in topological lattice models: Hamiltonian study of topological symmetry breaking*, Phys. Rev. B **84**, 125434 (2011).

[13] E. Witten, *Quantum field theory and the Jones polynomial*, Comm. Math. Phys. **121**, 351 (1989).

[14] G. Moore and N. Seiberg, *Taming the conformal zoo* Phys. Lett. B, **220**, 422 (1989).

[15] E. Lieb, T. Schultz, D. Mattis, *Two Soluble Models of an Antiferromagnetic Chain*, Ann. Phys. **16**, 407 (1961).

[16] R.J. Elliott, P. Pfeuty, C. Wood, *Ising model with a transverse field*, Phys. Rev. Lett. **25**, 443 (1970).

[17] P. Pfeuty, *The one-dimensional Ising model with a transverse field*, Ann. Phys. **57**, 79 (1970).

[18] P. Di Francesco, P. Mathieu, D. Sénéchal, *Conformal field theory*, Springer, New York (1999).

[19] D.S. Fisher, *Random antiferromagnetic quantum spin chains*, Phys. Rev. B **50**, 3799 (1994).

[20] T. Månsson, V. Lahtinen, J. Suorsa, E. Ardonne, in preparation.

[21] M. Oshikawa, I. Affleck, *Boundary conformal field theory approach to the critical two-dimensional Ising model with a defect line*, Nucl.Phys. B**495**, 533 (1997).